arXiv:1211.6809v1 [math.PR] 29 Nov 2012

A multiparameter Garsia-Rodemich-Rumsey inequality and some applications

Yaozhong Hu* and Khoa Le
Department of Mathematics, University of Kansas
405 Snow Hall, Lawrence, Kansas 66045-2142

Abstract

We extend the classical Garsia-Rodemich-Rumsey inequality to the multiparameter situation. The new inequality is applied to obtain some joint Hölder continuity along the rectangles for fractional Brownian fields $W(t,x)$ and for the solution $u(t,y)$ of stochastic heat equation with additive white noise.

1 Introduction

Let the function $\Psi : [0,\infty) \to [0,\infty)$ be non decreasing with $\lim\limits_{u\to\infty} \Psi(u) = \infty$ and let the function $p : [0,1] \to [0,1]$ be continuous and non decreasing with $p(0) = 0$. Set

$$\begin{cases} \Psi^{-1}(u) = \sup_{\Psi(v)\le u} v & \text{if } \Psi(0) \le u < \infty \\ \\ p^{-1}(u) = \max_{p(v)\le u} v & \text{if } 0 \le u \le p(1) \end{cases}$$

The celebrated Garsia-Rodemich-Rumsey inequality [6] takes the following form:

Lemma 1.1 *Let f be a continuous function on $[0,1]$ and suppose that*

$$\int_0^1 \int_0^1 \Psi\left(\frac{|f(x)-f(y)|}{p(x-y)}\right) dxdy \le B < \infty\,.$$

Then for all $s,t \in [0,1]$ we have

$$|f(s)-f(t)| \le 8 \int_0^{|s-t|} \Psi^{-1}\left(\frac{4B}{u^2}\right) dp(u). \tag{1.1}$$

This Garsia-Rodemich-Rumsey lemma 1.1 is very powerful in the study of the sample path Hölder continuity of a stochastic process and in other occasions. For example

---

[*] Y. Hu is partially supported by a grant from the Simons Foundation #209206.
Key words: Joint Hölder continuity; Garsia-Rodemich-Rumsey inequality; sample path property; Gaussian processes, fractional Brownian fields; stochastic heat equations with additive noises.
AMS subject classification (2010): 60G17, 26A16, 60G60, 60G15, 60H15.

if $\Psi(u) = |u|^p$ and $p(u) = |u|^{\alpha+1/p}$, where $p\alpha > 1$, the inequality (1.1) implies the following Sobolev imbedding inequality

$$|f(s) - f(t)| \le C_{\alpha,p}|t-s|^{\alpha-1/p}\left(\int_0^1\int_0^1 \frac{|f(x)-f(y)|^p}{|x-y|^{\alpha p+1}}dxdy\right)^{1/p}. \tag{1.2}$$

The Garsia-Rodemich-Rumsey lemma has been extended to several parameter or infinite many parameters. But the parameter space are assumed to have a distance (metric space) and the Garsia-Rodemich-Rumsey lemma is with respect to that distance. This method immediately yields the following result for a fractional Brownian field $W^H(x)$ of Hurst parameter $H = (H_1, \cdots, H_d)$, then for any β_i with $\beta_i < H_i$, $i = 1, \cdots, d$, one has

$$|W(y) - W(x)| \le L\sum_{i=1}^d |y_i - x_i|^{\beta_i}, \tag{1.3}$$

where L is an integrable random variable. One can improve this result (Remark 4.5) by our version of multiparameter Garsia-Rodemich-Rumsey inequality. We do not seek for a suitable metric but rather deal directly with the multidimensional nature of the parameter space.

Let us explain our motivation by considering the two parameter fractional Brownian field $\{W(x_1, x_2), (x_1, x_2) \in [0,1]^2\}$ of Hurst parameter $H = (H_1, H_2)$. Given two points x and y in $\mathbb{R}^2$, we consider the increment of W along with the rectangle determined by $x = (x_1, x_2)$ and $y = (y_1, y_2)$:

$$\square W := W(y_1, y_2) - W(x_1, y_2) - W(x_2, y_1) + W(x_1, x_2). \tag{1.4}$$

In [7], using a two-parameter version (1.2), the author showed that for any β_1, β_2 with $\beta_1 < H_1$ and $\beta_2 < H_2$, there is an integrable random constant L_{β_1,β_2} such that

$$|\square W| \le L_{\beta_1,\beta_2}|y_1 - x_1|^{\beta_1}|y_2 - x_2|^{\beta_2}. \tag{1.5}$$

The above result was also obtained in [1] based on a two-parameter version of Kolmogorov continuity theorem. Along the paper (in Corollary 4.4), we shall see that the following sharper inequality than (1.5) holds

$$|\square W| \le L_{H_1,H_2}|y_1 - x_1|^{H_1}|y_2 - x_2|^{H_2}\sqrt{|\log\left(|y_1 - x_1||y_2 - x_2|\right)|}. \tag{1.6}$$

Consequently, this estimate implies

$$\begin{aligned} |W(x_1, x_2) - W(y_1, y_2)| \le L_{H_1,H_2}\Big(&|x_1 - y_1|^{H_1}|x_2|^{H_2}\sqrt{|\log(|x_1 - y_1||x_2|)|} \\ &+ |x_1|^{H_1}|x_2 - y_2|^{H_2}\sqrt{|\log(|x_2 - y_2||x_1|)|}\Big) \end{aligned}$$

which improves (1.3). To our best knowledge, the estimate (1.6) is the first of its kind in literature. We shall call such property as in (1.6) or (1.5) joint Hölder continuity. It turns out that a large class of Gaussian fields enjoys sample path joint Hölder continuity (Theorem 4.3.)

Our method is first formulate and prove a multiparameter version of the classical Garsia-Rodemich-Rumsey inequality (1.1). The generalized inequality is then applied to obtain sample path joint Hölder continuity for random fields. Our result generalizes the results in [6], [7] and provides a different approach for sample path continuity problem of random fields (compare to the approach in [1], [2] and [9].)

The paper is structured as follows. In Section 2, we shall state and prove our multiparameter version of the Garsia-Rodemich-Rumsey lemma. The idea is to use induction on the dimension of the parameter space after some observations of the property of operator $\Box$ defined by (1.4). Some part of the proof is similar to the original proof of Garsia-Rodemich-Rumsey [6] with some modification. However, we feel it is more appropriate to give a detailed proof.

In Section 3, we introduce a multiparameter version of Kolmogorov continuity criteria (Theorem 3.1). To our best knowledge, a two-parameter of Theorem 3.1 first appeared in [1].

Section 4 is devoted for the study of sample path continuity for Gaussian fields. We give a sufficient condition for a Gaussian field to possess sample path continuity (Theorem 4.3). We also derive the estimate (1.6) for fractional Gaussian field. In Section 5, we shall study the joint Hölder continuity of solution of a stochastic heat equation with additive space-time white noise.

2 Multiparameter Garsia-Rodemich-Rumsey inequality

We state the following technical lemma which generalizes a crucial argument used in [6] in the proof of Lemma 1.1.

Lemma 2.1 *Let $(\Omega, \mathcal{F})$ be a measurable space and let μ be a positive measure on $(\Omega, \mathcal{F})$. Let $g : \Omega \times [0,1] \to \mathbb{R}^m$ be a measurable function such that*

$$\int_0^1 \int_0^1 \int_\Omega \Psi\left(\frac{|g(z,t)-g(z,s)|}{p(|t-s|)}\right)\mu(dz)dsdt \le B < \infty .$$

Then there exist two decreasing sequences $\{t_k, k = 0, 1, \cdots\}$ and $\{d_k, k = 0, 1, \cdots\}$ with

$$t_k \le d_{k-1} = p^{-1}\left(\frac{1}{2}p(t_{k-1})\right), \quad k = 1, 2, \cdots \tag{2.1}$$

such that the following inequality holds

$$\int_\Omega \Psi\left(\frac{|g(z,t_k)-g(z,t_{k-1})|}{p(|s-t|)}\right)\mu(dz) \le \frac{4B}{d_{k-1}^2} . \tag{2.2}$$

Proof We follow the argument in [6]. Let

$$I(t) = \int_0^1 \int_\Omega \Psi\left(\frac{|g(z,t)-g(z,s)|}{p(|t_k - t_{k-1}|)}\right)\mu(dz)ds.$$

From the assumption $\int_0^1 I(t)dt \le B$ it follows that there is some $t_0 \in (0,1)$ such that

$$I(t_0) \le B.$$

Now we can describe how to obtain the sequences d_k and t_k recursively for $k = 1, 2, \cdots$. Given t_{k-1}, define

$$d_{k-1} = p^{-1}\left(\frac{1}{2}p(t_{k-1})\right) .$$

Then we choose $t_k \leq d_{k-1}$ such that

$$I(t_k) \leq \frac{2B}{d_{k-1}} \tag{2.3}$$

and

$$\int_\Omega \Psi\left(\frac{|g(z,t_k)-g(z,t_{k-1})|}{p(|t_k-t_{k-1}|)}\right)\mu(dz) \leq \frac{2I(t_{k-1})}{d_{k-1}}\,. \tag{2.4}$$

It is always possible to find t_k such that the inequalities (2.3) and (2.4) hold simultaneously, since each of the two inequalities can be violated only on a set of t_k's of measure strictly less than $\frac{1}{2}d_{k-1}$. Now (2.3) and (2.4) gives

$$\int_\Omega \Psi\left(\frac{|g(z,t_k)-g(z,t_{k-1})|}{p(|t_k-t_{k-1}|)}\right)\mu(dz) \leq \frac{2I(t_{k-1})}{d_{k-1}} \leq \frac{4B}{d_{k-1}d_{k-2}} \leq \frac{4B}{d_{k-1}^2}\,.$$

This is (2.2). ■

Let $x=(x_1,\ldots,x_n)$ and $y=(y_1,\ldots,y_n)$ be in $\mathbb{R}^n$. We denote $x'=(x_1,\ldots,x_{n-1})$ and $y'=(y_1,\ldots,y_{n-1})$. For each integer $k=1,2,\cdots,n$, we define

$$V_{k,y}x=(x_1,\ldots,x_{k-1},y_k,x_{k+1},\ldots,x_n).$$

Let f be a function from $\mathbb{R}^n$ to $\mathbb{R}^m$. We define the operator $V_{k,y}$ acting on f in the following way:

$$V_{k,y}f(x)=f(V_{k,y}x)$$

or $V_{k,y}f=f\circ V_{k,y}$ in short. It is straight forward to verify that

$$V_{k,y}V_{k,y}=V_{k,y}$$

and

$$V_{k,y}V_{l,y}=V_{l,y}V_{k,y}$$

for $k\neq l$. Next, we define the joint increment of a function f on an n-dimensional rectangle, namely

$$\Box_y^n f(x)=\prod_{k=1}^n (I-V_{k,y})f(x)$$

where I denotes the identity operator.

Example 2.2 *If $n=2$, then it is easy to see that $\Box_y^2 f(x)=f(y_1,y_2)-f(x_1,y_2)-f(y_1,x_2)+f(x_1,y_2)$, which is the increment of f over the rectangle containing the two points x and y with all sides parallel to the axis. In particular, if $f(x_1,x_2)=x_1x_2$, then $\Box_y^2 f(x)=(x_1-y_1)(x_2-y_2)$, which is the area of the rectangle. In a more general case, when f has the form $f(x)=\prod_{j=1}^n f_j(x_j)$, then*

$$\Box_y^n f(x)=\prod_{j=1}^n [f_j(x_j)-f_j(y_j)]\,.$$

The following simple identity enable us to show our theorem by induction and plays an essential role in our approach:

$$\begin{aligned}\Box_y^n f(x) &= \prod_{k=1}^{n-1}(I - V_{k,y})f(x) - V_{n,y}\prod_{k=1}^{n-1}(I - V_{k,y})f(x) \\ &= \Box_{y'}^{n-1} f(x', x_n) - \Box_{y'}^{n-1} f(x', y_n). \end{aligned} \tag{2.5}$$

We are now in the position to state our general version of Lemma 1.1.

Theorem 2.3 *Let $f(x)$ be a continuous function on $[0,1]^n$ and suppose that*

$$\int_{[0,1]^n}\int_{[0,1]^n} \Psi\left(\frac{|\Box_y^n f(x)|}{\prod_{k=1}^n p_k(|x_k - y_k|)}\right) dxdy \le B < \infty\,. \tag{2.6}$$

Then for all $s, t \in [0,1]^n$ we have

$$|\Box_s^n f(t)| \le 8^n \int_0^{|s_1-t_1|}\cdots\int_0^{|s_n-t_n|} \Psi^{-1}\left(\frac{4^n B}{u_1^2\cdots u_n^2}\right) dp_1(u_1)\cdots dp_n(u_n). \tag{2.7}$$

Proof We proceed by induction on n. For $n = 1$, it coincides with the original Garsia-Rodemich-Rumsey inequality (1.1). Suppose (2.7) holds for $n-1$. Let f be a continuous function on $[0,1]^n$. For any $x', y' \in \mathbb{R}^{n-1}$ and any $s \in [0,1]$, put

$$g(x', y', s) = \frac{\Box_{y'}^{n-1} f(x', s)}{\prod_{k=1}^{n-1} p_k(|x'_k - y'_k|)}.$$

Let $\Omega = [0,1]^{n-1} \times [0,1]^{n-1}$, $z = (x', y')$. By (2.5) we can rewrite (2.6) as

$$\int_0^1\int_0^1\int_\Omega \Psi\left(\frac{|g(z,s) - g(z,t)|}{p_n(|s-t|)}\right) dzdsdt \le B < \infty.$$

Applying Lemma 2.1, we can find sequences $\{t_k\}$ and $\{d_k\}$ such that

$$t_k \le d_{k-1} = p_n^{-1}\left(\frac{1}{2}p_n(t_{k-1})\right) \tag{2.8}$$

and

$$\int_\Omega \Psi\left(\frac{|g(z,t_k) - g(z,t_{k-1})|}{p_n(t_k - t_{k-1})}\right) dz \le \frac{4B}{d_{k-1}^2}. \tag{2.9}$$

For each $k \in \mathbb{N}$ and $x' \in [0,1]^{n-1}$, let

$$h_k(x') = \frac{f(x', t_k) - f(x', t_{k-1})}{p_n(|t_k - t_{k-1}|)}.$$

Again from (2.5) it follows

$$\begin{aligned}\frac{\Box_{y'}^{n-1} h_k(x')}{\prod_{i=1}^{n-1} p_i(|x'_i - y'_i|)} &= \frac{\Box_{y'}^{n-1} f(x', t_k) - \Box_{y'}^{n-1} f(x', t_{k-1})}{\prod_{i=1}^{n-1} p_i(|x'_i - y'_i|)p_n(|t_k - t_{k-1}|)} \\ &= \frac{g(x', y', t_k) - g(x', y', t_{k-1})}{p_n(|t_k - t_{k-1}|)}.\end{aligned}$$

Thus, the inequality (2.9) becomes

$$\int_{[0,1]^{n-1}}\int_{[0,1]^{n-1}}\Psi\left(\frac{\square_{y'}^{n-1}h_k(x')}{\prod_{i=1}^{n-1}p_i(|x_i'-y_i'|)}\right)dx'dy'\le\frac{4B}{d_{k-1}^2}.$$

Now, by our induction hypothesis, for every $k\ge 1$, $a,b\in[0,1]^{n-1}$, $a=(a_1,\dots,a_{n-1})$ and $b=(b_1,\dots,b_{n-1})$,

$$\begin{aligned}|\square_a^{n-1}h_k(b)| &\le 8^{n-1}\int_0^{|a_1-b_1|}\cdots\int_0^{|a_{n-1}-b_{n-1}|}\Psi^{-1}\left(\frac{4^nB}{u_1^2\cdots u_{n-1}^2d_{k-1}^2}\right)\\ &\qquad dp_1(u_1)\cdots dp_{n-1}(u_{n-1}).\end{aligned}$$

Denoting $A=[0,|a_1-b_1|]\times\cdots\times[0,|a_{n-1}-b_{n-1}|]$ and $dp(u_1,\cdots,u_{n-1})=dp_1(u_1)\cdots dp_{n-1}(u_{n-1})$, the above inequality can be rewritten as

$$|\square_a^{n-1}f(b,t_k)-\square_a^{n-1}f(b,t_{k-1})|\le 8^{n-1}\int_A\Psi^{-1}\left(\frac{4^nB}{u_1^2\cdots u_{n-1}^2d_{k-1}^2}\right)dp(u_1,\cdots,u_{n-1})p_n(t_{k-1}-t_k). \tag{2.10}$$

On the other hand, by (2.8), we have

$$\begin{aligned}p_n(t_{k-1}-t_k) &\le p_n(t_{k-1})\\ &= 2p(d_{k-1})\le 4\left[p(d_{k-1})-p(d_k)\right].\end{aligned}$$

Combining this inequality with (2.10) yields

$$\begin{aligned}&|\square_a^{n-1}f(b,t_0)-\square_a^{n-1}f(b,0)|\\ \le&\sum_{k=1}^{\infty}|\square_a^{n-1}f(b,t_k)-\square_a^{n-1}f(b,t_{k-1})|\\ \le& 8^{n-1}\sum_{k=1}^{\infty}4\left[p(d_{k-1})-p(d_k)\right]\int_A\Psi^{-1}\left(\frac{4^nB}{u_1^2\cdots u_{n-1}^2d_{k-1}^2}\right)dp(u_1,\cdots,u_{n-1})\\ \le& 8^{n-1}\sum_{k=1}^{\infty}4\int_{d_k}^{d_{k-1}}\int_A\Psi^{-1}\left(\frac{4^nB}{u_1^2\cdots u_{n-1}^2u_n^2}\right)dp(u_1,\cdots,u_{n-1})dp_n(u_n)\\ \le& 8^{n-1}4\int_0^1\int_A\Psi^{-1}\left(\frac{4^nB}{u_1^2\cdots u_n^2}\right)dp(u_1,\cdots,u_{n-1})dp_n(u_n).\end{aligned}$$

With $f(x',1-x_n)$ replaced $f(x',x_n)$ we can obtain the same bound for

$$|\square_a^{n-1}f(b,t_0)-\square_a^{n-1}f(b,1)|.$$

Hence, for every $a,b\in[0,1]^{n-1}$,

$$|\square_a^{n-1}f(b,1)-\square_a^{n-1}f(b,0)|\le 8^n\int_0^1\int_A\Psi^{-1}\left(\frac{4^nB}{u_1^2\cdots u_n^2}\right)dp(u_1,\cdots,u_{n-1})dp_n(u_n). \tag{2.11}$$

To obtain (2.7) for general s,t in $[0,1]^n$, we set

$$\bar f(t',\tau)=f(t',s_n+\tau(t_n-s_n))\text{ for }\tau\in[0,1]$$

and

$$\bar{p}_n(u) = p_n(u|s_n - t_n|).$$

Upon restricting the range of the integration in (2.6) and carrying out a change of variables we get

$$\int_{[0,1]^n} \int_{[0,1]^n} \Psi\left(\frac{|\Box_y^n \bar{f}(x)|}{\prod_{k=1}^{n-1} p_k(|x_k - y_k|)\bar{p}_n(|x_n - y_n|)} \right) dxdy \leq \frac{B}{|s_n - t_n|^2}.$$

Thus, by (2.11), we deduce

$$\begin{aligned} |\Box_s^n f(t)| &= |\Box_{s'}^{n-1} \bar{f}(t', 1) - \Box_{s'}^{n-1} \bar{f}(t', 0)| \\ &\leq 8^n \int_0^1 \int_0^{|s_1 - t_1|} \cdots \int_0^{|s_{n-1} - t_{n-1}|} \Psi^{-1}\left(\frac{4^n B}{u_1^2 \cdots u_n^2 |s_n - t_n|^2} \right) \\ &\qquad dp(u_1, \cdots, u_{n-1}) dp_n(u_n|s_n - t_n|). \end{aligned}$$

Another change of variables yields (2.7). ■

3 Sample path Hölder continuity of random fields

In this section, given a random field W, we study sample path continuity property. The first application of Theorem 2.3 is the following criteria for joint continuity of sample paths which is similar to Kolmogorov continuity theorem, which we shall call joint Kolmogorov continuity theorem.

Theorem 3.1 *Let W be a random field on $\mathbb{R}^n$. Suppose there exist positive constants α, β_k $(1 \leq k \leq n)$ and K such that for every x, y in $[0,1]^n$,*

$$\mathbb{E}\left[\left| \Box_y^n W(x) \right|^\alpha \right] \leq K \prod_{k=1}^n |x_k - y_k|^{1+\beta_k}.$$

Then, for every $\epsilon = (\epsilon_1, \ldots, \epsilon_n)$ with $0 < \epsilon_k \alpha < \beta_k$ $(1 \leq k \leq n)$, there exist a random variable η with $\mathbb{E}\eta^\alpha \leq K$, such that the following inequality holds almost surely

$$|\Box_t^n W(s)| \leq C\eta(\omega) \prod_{k=1}^n |t_k - s_k|^{\beta_k \alpha^{-1} - \epsilon_k}$$

for all s, t in $[0,1]^n$, where C is a constant defined by

$$C = 8^n 4^{n/\alpha} \prod_{k=1}^n \left(1 + \frac{2}{\beta_k - \alpha\epsilon_k} \right).$$

Proof Let $\Psi(u) = |u|^\alpha$, $p_k(u) = |u|^{\gamma_k}$ where $\gamma_k \in (\frac{2}{\alpha}, \frac{2+\beta_k}{\alpha})$, $1 \leq k \leq n$. A direct application of Theorem 2.3 gives that for all s, t in $[0,1]^n$

$$|\Box_s^n W(t)| \leq 8^n \prod_{k=1}^n \frac{\gamma_k |t_k - s_k|^{\gamma_k - \frac{2}{\alpha}}}{\gamma_k - \frac{2}{\alpha}} \left(4^n \iint_{[0,1]^{2n}} \frac{|\Box_y^n W(x)|^\alpha}{\prod_{k=1}^n |x_k - y_k|^{\alpha\gamma_k}} dxdy \right)^{\frac{1}{\alpha}}. \tag{3.1}$$

Let

$$B(\omega) = \iint_{[0,1]^{2n}} \Psi\left(\frac{|\Box_y^n W(x)|}{\prod_{k=1}^n p_k(x_k - y_k)}\right) dxdy.$$

From our assumption and Fubini-Tonelli's theorem,

$$\begin{aligned}
\mathbb{E}B &= \iint_{[0,1]^{2n}} \frac{\mathbb{E}|\Box_y^n W(x)|^\alpha}{\prod_{k=1}^n |x_k - y_k|^{\alpha\gamma_k}} dxdy \\
&\leq K \iint_{[0,1]^{2n}} \prod_{k=1}^n |x_k - y_k|^{1+\beta_k - \alpha\gamma_k} dxdy < \infty.
\end{aligned}$$

Hence, the event $\Omega^* = \{\omega : B(\omega) < \infty\}$ has probability one. Therefore for each ω in Ω^*, the inequality (3.1) gives

$$|\Box_s^n W(t,\omega)| \leq 8^n \prod_{k=1}^n \frac{\gamma_k |t_k - s_k|^{\gamma_k - \frac{2}{\alpha}}}{\gamma_k - \frac{2}{\alpha}} (4^n B(\omega))^{\frac{1}{\alpha}}$$

for every s, t in $[0,1]^n$. For each k, the power $\gamma_k - \frac{2}{\alpha}$ can be made arbitrarily close to $\frac{\beta_k}{\alpha}$. This completes the proof with $\eta = B^{1/\alpha}$. ■

Remark 3.2 *The result obtained by Ral'chenko [7] was the inequality (3.1) in the case* $n = 2$.

4 Sample path continuity of Gaussian fields

We now focus on sample path continuity of Gaussian random fields. In case of Gaussian processes ($n = 1$), one of the first sufficient and necessary conditions for sample path continuity was given by Fernique [3] (see also [5]). Namely, let $p(u)$ be an increasing positive function such that

$$\mathbb{E}|W(x) - W(y)|^2 \leq p^2(|x - y|)$$

for any pair (x, y) in $[0,1]^2$. Then Fernique [3] showed that a sufficient condition for almost sure continuity of the process $(W(x), 0 \leq x \leq 1)$ is

$$\int_0^1 \frac{p(u)}{u\sqrt{\log \frac{1}{u}}} du < \infty.$$

In the original paper of Garsia-Rodemich-Rumsey [6], the authors also observed that the above condition is equivalent to the condition (by integration by part)

$$\int_0^1 \sqrt{\log \frac{1}{u}} dp(u) < \infty.$$

Later, it was shown that the above condition is also necessary [4, 8]. In case of Gaussian fields, recent progress on modulus of continuity of Gaussian random fields has been reported in [9, 2].

Let W be a centered Gaussian random field with covariance function

$$\mathbb{E}\left[W(x)W(y)\right] = Q(x,y). \tag{4.1}$$

We will always assume that Q is a continuous function of x and y. For any fixed x, y, the random variable $\Box_y^n W(x)$ is also Gaussian with mean zero. In the following proposition, we compute its variance.

Proposition 4.1 *Let W be a centered Gaussian random field with covariance function given by (4.1). Then*

$$\mathbb{E}\left[|\Box_y^n W(x)|^2\right] = \Box_{(y,y)}^{2n} Q(x,x). \tag{4.2}$$

Furthermore, if the covariance function Q has the following product form

$$Q(x,y) = \prod_{k=1}^{n} Q_k(x_k, y_k) \tag{4.3}$$

then (4.2) is simplified as

$$\mathbb{E}\left[|\Box_y^n W(x)|^2\right] = \prod_{k=1}^{n} \left[Q_k(x_k,x_k) - Q_k(x_k,y_k) - Q_k(y_k,x_k) + Q_k(y_k,y_k)\right]. \tag{4.4}$$

Proof We calculate the variance directly as follows

$$\begin{aligned}
\mathbb{E}\left[|\Box_y^n W(x)|^2\right] &= \mathbb{E}\left[\Box_y^n W(x) \cdot \Box_y^n W(x)\right] \\
&= \mathbb{E}\left[\Box_{(y,y)}^{2n} W(x)W(x)\right] \\
&= \Box_{(y,y)}^{2n} \mathbb{E}\left[W(x)W(x)\right] \\
&= \Box_{(y,y)}^{2n} Q(x,x).
\end{aligned}$$

The identity (4.2) follows. To prove (4.4), we notice that the pair of operators $(I - V_{k,(y,y)})(I - V_{n+k,(y,y)})$ transforms the k-th factor of Q in (4.3) to

$$Q_k(x_k,x_k) - Q_k(x_k,y_k) - Q_k(y_k,x_k) + Q_k(y_k,y_k).$$

Since the operators $I - V_{k,(y,y)}$, $(1 \le k \le 2n)$ are commutative, we can write

$$\begin{aligned}
\Box_{(y,y)}^{2n} Q(x,x) &= \prod_{k=1}^{n} (I - V_{k,(y,y)})(I - V_{n+k,(y,y)}) Q(x,x) \\
&= \prod_{k=1}^{n} \left[Q_k(x_k,x_k) - Q_k(x_k,y_k) - Q_k(y_k,x_k) + Q_k(y_k,y_k)\right].
\end{aligned}$$

Hence, the identity (4.4) follows. ■

Definition 4.2 *Let f be a continuous function on $\mathbb{R}^n$. We call a set of non-negative even functions $\{p_1, \ldots, p_n\}$ joint modulus of continuity of f if*

(i) For each $1 \le k \le n$, $p_k(0) = 0$, and p_k is non-decreasing and continuous.

(ii) For every pair (s,t) in $\mathbb{R}^{2n}$, the following inequality holds

$$|\Box_s^m f(t)| \leq \prod_{k=1}^{n} p_k(|t_k - s_k|)\,.$$

In view of Theorem 2.3 and Theorem 3.1, the continuity of sample paths is governed by the joint modulus of continuity of $\Box^{2n}_{(y,y)} Q(x,x)$. Such modulus of continuity always exists. For instance, we can define a joint modulus of continuity for $\Box^{2n}_{(y,y)} Q(x,x)$ as follows. We set

$$p_1(u) = \sup_{x,y\in[0,1]^n:|x_1-y_1|\leq u} \left[\Box^{2n}_{(y,y)} Q(x,x)\right]^{\frac{1}{2}}.$$

Given $p_1, \dots, p_{k-1}$, define

$$p_k(u) = \sup_{x,y\in[0,1]^n:|x_k-y_k|\leq u} \frac{\left[\Box^{2n}_{(y,y)} Q(x,x)\right]^{\frac{1}{2}}}{\prod_{j=1}^{k-1} p_j(|x_j-y_j|)},$$

in which we have adopted the convention $0/0=0$. It follows immediately that p_k's are non-decreasing and continuous. Furthermore, we have $p_k(0)=0$ and

$$\Box^{2n}_{(y,y)} Q(x,x) \leq \prod_{k=1}^{n} p_k^2(|x_k - y_k|)\,. \tag{4.5}$$

Namely, $\{p_1,p_1,p_2,p_2,\dots,p_n,p_n\}$ is a modulus of continuity for $\Box^{2n}_{(y,y)} Q(x,x)$. We also call $\{p_1,\cdots,p_n\}$ a modulus of continuity for $\Box^{2n}_{(y,y)} Q(x,x)$.

In the following theorem, we give a sufficient condition for almost sure continuity of a Gaussian random field.

Theorem 4.3 *Let W be a centered Gaussian random field with covariance function given by (4.1), and p_k $(1\leq k\leq n)$ be a modulus of continuity for $\Box^{2n}_{(y,y)} Q(x,x)$, namely the inequality (4.5) is satisfied. Suppose that*

$$\sum_{k=1}^{n} \int_0^1 \left(\log\frac{1}{u}\right)^{\frac{1}{2}} dp_k(u) < \infty. \tag{4.6}$$

Then, with probability one W has continuous sample path. Furthermore, there is a random variable $B(\omega)$, with finite expectation, such that

$$|\Box_y^n W(x)| \leq 2\cdot 8^n \int_0^{|x_1-y_1|} \cdots \int_0^{|x_n-y_n|} \left(\log\frac{4^n B(\omega)}{u_1^2\cdots u_n^2}\right)^{\frac{1}{2}} dp_1(u_1)\cdots dp_n(u_n). \tag{4.7}$$

Proof We set $\Psi(x) = e^{x^2/4}$ and

$$B(\omega) = \iint_{[0,1]^{2n}} \exp\left[\frac{|\Box_y^n W(x)|^2}{4\prod_{k=1}^n p_k^2(|x_k-y_k|)}\right] dxdy.$$

Lemma 2.3 immediately gives us (4.7) for ω such that $B(\omega)$ is finite. To see (4.7) indeed holds for almost every ω, it is sufficient to show that B has finite expectation. We notice that the random variable

$$N=\frac{\square_y^n W(x)}{\prod_{k=1}^n p_k(|x_k-y_k|)}$$

is Gaussian, has mean zero and variance less than or equal to one. Thus, an application of Stirling's formula gives

$$\begin{aligned}\mathbb{E}\exp\left(\frac{N^2}{4}\right) &= \sum_{k=0}^{\infty}\frac{\mathbb{E}N^{2k}}{4^k k!}\\ &= 1+\sum_{k=1}^{\infty}\frac{(2k)!}{8^k\,(k!)^2}(\mathbb{E}N^2)^k\\ &\le 1+\frac{1}{2}\sum_{k=1}^{\infty}8^{-k}=\frac{15}{14}.\end{aligned}$$

Hence

$$\mathbb{E}B=\iint_{[0,1]^{2n}}\mathbb{E}\exp\left(\frac{N^2}{4}\right)dxdy\le\frac{15}{14}$$

and the proof is complete. ■

As an application of the above theorem, we obtain joint continuity for sample paths of fractional Brownian field, as mentioned in (1.6).

Corollary 4.4 *Let W^H be a fractional Brownian field on $\mathbb{R}^n$ with Hurst parameter $H=(H_1,\dots,H_n)$. Then, there exist an integrable random constant $C_{n,H_1,\cdots,H_n}$ such that for every x,y in $[0,1/2]^n$*

$$|\square_y^n W^H(x)|\le C_{n,H_1,\cdots,H_n}\sqrt{\left|\log\left(\prod_{k=1}^n|x_k-y_k|\right)\right|}\;\prod_{k=1}^n|x_k-y_k|^{H_k}\,. \tag{4.8}$$

Proof The covariance function of a fractional Brownian field is given by

$$\mathbb{E}\left[W^H(x)W^H(y)\right]=\prod_{k=1}^n R_k(x_k,y_k),$$

where

$$R_k(s,t)=\frac{1}{2}\left[|s|^{2H_k}+|t|^{2H_k}-|s-t|^{2H_k}\right]\,,\quad\forall\, s,t\in\mathbb{R}\,.$$

By Proposition 4.1, we obtain the second moment for $\square_y^n W^H(x)$

$$\mathbb{E}|\square_y^n W^H(x)|^2=\prod_{k=1}^n|x_k-y_k|^{2H_k}.$$

This means that $p_i(u)=u^{H_i}\,, i=1,2,\cdots,n$ are the modulus of continuity of $\square_{(y,y)}^{2n}Q(x,x)$. From Theorem 4.3 it follows that

$$|\square_y^n W(x)|\le C_{n,H_1,\cdots,H_n}\int_0^{|x_1-y_1|}\cdots\int_0^{|x_n-y_n|}\sqrt{\log\frac{1}{u_1\cdots u_n}}\,u_1^{H_1-1}\cdots u_n^{H_n-1}du_1\cdots du_n.$$

Now an elementary computation yields the inequality (4.8). ■

Remark 4.5 *In [2], it is shown that for fractional Brownian field W^H on $\mathbb{R}^n$, there exist a random variable $A_1 = A_1(\omega) > 0$ of finite moments of any order and an event Ω_1^* of probability 1 such that for any $\omega \in \Omega_1^*$,*

$$\sup_{s,t\in[0,1]^n} \frac{|W^H(s,\omega) - W^H(t,\omega)|}{\sum_{j=1}^n |s_j - t_j|^{H_j} \sqrt{\log\left(3 + |s_j - t_j|^{-1}\right)}} \le A_1(\omega)\,.$$

Our method leads to a slightly stronger estimate. Indeed, for every s,t in $[0,1/2]^n$, with s and $(0,0,\ldots,0,t_n)$, the estimate (4.8) gives the increment along an edge of the n-dimensional rectangle $[s_1,t_1]\times\cdots\times[s_n,t_n]$

$$\begin{aligned}&\left|W^H(s_1,\cdots,s_n) - W^H(s_1,\cdots,s_{n-1},t_n)\right|\\ &\qquad \le A_1(\omega)\left(\prod_{k=1}^{n-1} s_k^{H_k}\right)\left|\log\prod_{k=1}^{n-1} s_k\right|^{1/2} |s_n - t_n|^{H_n}\left|\log|s_n - t_n|\right|^{1/2}.\end{aligned}$$

Similarly, we can obtain analogue estimates along any edge of the n-dimensional rectangle $[s_1,t_1]\times\cdots\times[s_n,t_n]$. The increment along the diagonal is majorized by the total increments along all the edges connecting s and t. Hence, this argument yields the following estimate

$$\left|W^H(s) - W^H(t)\right| \le A_1(\omega)\sum_{k=1}^n\left(\prod_{j\ne k} s_j^{H_j}\right)\left|\log\prod_{j\ne k} s_j\right|^{1/2} |s_k - t_k|^{H_k}\left|\log|s_k - t_k|\right|^{1/2}. \tag{4.9}$$

5 Stochastic heat equations with additive space time white noise

In this section let us consider the following one dimensional stochastic differential equation

$$\begin{cases} \frac{\partial u}{\partial t} = \frac{1}{2}\Delta u + \dot W & 0 < t \le T\,,\ y\in\mathbb{R} \\ u(0,y) = 0 & y \in \mathbb{R}\,, \end{cases} \tag{5.1}$$

where $\Delta u = \frac{\partial^2}{\partial y^2}u$, W is space time standard Brownian sheet, and $\dot W = \frac{\partial^2}{\partial t\partial y}W$. Let $p_t(y) = \dfrac{1}{\sqrt{2\pi t}}e^{-\frac{y^2}{2t}}$. Then the (mild) solution of the above equation is given by

$$u(t,y) = \int_0^t\int_{\mathbb{R}} p_{t-r}(y-z)W(dr,dz)\,,$$

where the above integral is the usual (Itô) stochastic integral (however, the integrand is simple. It is a deterministic function). The solution $u(t,y)$ is a Gaussian random field. It is known that $u(t,y)$ is Hölder continuous of exponent $\frac14-$ for time parameter and $\frac12-$ for space parameter. Namely, for any $\alpha < 1/4$ and any $\beta < 1/2$, there is a random constant $C_{\alpha,\beta}$ such that

$$|u(t,y) - u(s,x)| \le C_{\alpha,\beta}\left(|t-s|^\alpha + |x-y|^\beta\right)\,. \tag{5.2}$$

We are interested in the joint Hölder continuity of the solution $u(t,y)$. We need the following simple technical lemma.

Lemma 5.1 *Let a, b, δ be some positive numbers, where $a < b$, and let I, J be the integrations*

$$I = \int_a^b \frac{1}{\sqrt{r}} \left(1 - e^{-\frac{\delta^2}{2r}}\right) dr,$$

$$J = \int_0^a \frac{1}{\sqrt{r}} \left(1 - e^{-\frac{\delta^2}{2r}}\right) dr.$$

Then for every $\alpha \in [0, 1/2]$,

$$2(\sqrt{b} - \sqrt{a}) \left(1 - e^{-\frac{\delta^2}{2b}}\right) \le I \le 2(\sqrt{b} - \sqrt{a}) \left(1 - e^{-\frac{\delta^2}{2a}}\right)$$

and

$$J \le c_\alpha \delta^{2\alpha} a^{1/2-\alpha}.$$

Proof On the interval $a \le r \le b$, we have $1 - e^{-\frac{\delta^2}{2b}} \le 1 - e^{-\frac{\delta^2}{2r}} \le 1 - e^{-\frac{\delta^2}{2a}}$. The estimate for I is then a straightforward consequence. To estimate J, we first use integration by part to obtain

$$\begin{aligned} J &= 2\sqrt{r} \left(1 - e^{-\frac{\delta^2}{2r}}\right)_{r=0}^{r=a} + \int_0^a \left(\frac{d}{dr} e^{-\frac{\delta^2}{2r}}\right) 2\sqrt{r} dr \\ &= 2\sqrt{a} \left(1 - e^{-\frac{\delta^2}{2a}}\right) + \delta^2 \int_0^a e^{-\frac{\delta^2}{2r}} r^{-3/2} dr. \end{aligned}$$

By a change of variable $x = \frac{\delta}{\sqrt{2r}}$, we see that

$$J = 2\sqrt{a} \left(1 - e^{-\frac{\delta^2}{2a}}\right) + 2\sqrt{2}\delta \int_{\frac{\delta}{\sqrt{2a}}}^{\infty} e^{-x^2} dx.$$

If $\frac{\delta}{\sqrt{2a}} \ge 1$, since $\lim_{t\to\infty} \frac{\int_t^\infty e^{-x^2} dx}{(1-e^{-t^2})t^{-1}} = 0$, J is majorized by

$$J \le c\sqrt{a} \left(1 - e^{-\frac{\delta^2}{2a}}\right).$$

If $\frac{\delta}{\sqrt{2a}} \le 1$, the integration $\int_{\frac{\delta}{\sqrt{2a}}}^{\infty} e^{-x^2} dx$ is bounded by $\sqrt{\pi}/2$, thus J is majorized by

$$J \le 2\sqrt{a} \left(1 - e^{-\frac{\delta^2}{2a}}\right) + c\delta.$$

Therefore, for any $0 \le \alpha \le 1/2$, employing the elementary inequality $1 - e^{-x} \le c_\alpha x^\alpha$, we obtain

$$J \le c_\alpha \delta^{2\alpha} a^{1/2-\alpha}$$

and the lemma follows. ■

Theorem 5.2 *For every α in $[0, 1/4]$, there is an integrable random constant C_α such that for all $(s,x), (t,y)$ in $[0,1/2]^2$*

$$|u(t,y) - u(t,x) - u(s,y) + u(s,x)| \le C_\alpha |t-s|^{1/4-\alpha} |x-y|^{2\alpha} \sqrt{|\log(|t-s||x-y|)|}. \quad (5.3)$$

Proof $u(t,y)$ is a mean zero Gaussian field. The covariance of $u(t,y)$ and $u(s,x)$ is given by

$$\begin{aligned}\mathbb{E}[u(s,x)u(t,y)] &= \int_{\mathbb{R}^2} \chi_{[0,s]}(r)\chi_{[0,t]}(r)p_{s-r}(x-z)p_{t-r}(y-z)drdz \\ &= \int_{\mathbb{R}^2} f(s,x)f(t,y)drdz\,,\end{aligned}$$

where $f(s,x) = \chi_{[0,s]}(r)p_{s-r}(x-z)$.

We calculate the second moment of $\Box^2_{(s,x)}u(t,y)$ as follows

$$\begin{aligned}\mathbb{E}\left[\Box^2_{(s,x)}u(t,y)\right]^2 &= \mathbb{E}\Box^2_{(s,x)}u(t,y)\Box^2_{(s,x)}u(t,y) \\ &= \mathbb{E}\Box^4_{(s,x,s,x)}u(t,y)u(t,y) \\ &= \Box^4_{(s,x,s,x)}\mathbb{E}\left[u(t,y)u(t,y)\right] \\ &= \Box^4_{(s,x,s,x)}\int_{\mathbb{R}^2} f(t,y)^2 drdz \\ &= \int_{\mathbb{R}^2} \Box^4_{(s,x,s,x)}\left[f(t,y)f(t,y)\right]drdz \\ &= \int_{\mathbb{R}^2} \left[\Box^2_{(s,x)}f(t,y)\right]\left[\Box^2_{(s,x)}f(t,y)\right]drdz \\ &= \int_{\mathbb{R}^2} \left[\Box^2_{(s,x)}f(t,y)\right]^2 drdz\,,\end{aligned}$$

where

$$\begin{aligned}\left[\Box^2_{(s,x)}f(t,y)\right]^2 &= \left[f(s,x)-f(t,x)-f(s,y)+f(t,y)\right]^2 \\ &= f(s,x)^2+f(t,x)^2+f(s,y)^2+f(t,y)^2 \\ &\quad -2f(s,x)f(t,x)-2f(s,x)f(s,y)+2f(s,x)f(t,y) \\ &\quad +2f(t,x)f(s,y)-2f(t,x)f(t,y)-2f(s,y)f(t,y).\end{aligned}$$

Taking the integration with respect to z and using the following identity

$$\int_{\mathbb{R}} p_a(z-x)p_b(z-y)dz = p_{a+b}(x-y)$$

we obtain

$$\begin{aligned}&\mathbb{E}\left[\Box^2_{(s,x)}u(t,y)\right]^2 \\ =\ & \int_{\mathbb{R}}\left[2\chi_{[0,s]}(r)p_{2s-2r}(0)+2\chi_{[0,t]}(r)p_{2t-2r}(0)\right]dr \\ & +\int_{\mathbb{R}}\left[-2\chi_{[0,s\wedge t]}p_{s+t-2r}(0)-2\chi_{[0,s]}p_{2s-2r}(x-y)+2\chi_{[0,s\wedge t]}p_{s+t-2r}(x-y)\right]dr \\ & +\int_{\mathbb{R}}\left[2\chi_{[0,s\wedge t]}p_{s+t-2r}(x-y)-2\chi_{[0,t]}p_{2t-2r}(x-y)-2\chi_{[0,s\wedge t]}p_{s+t-2r}(0)\right]dr \\ =\ & 2\int_0^s\left[p_{2s-2r}(0)-p_{2s-2r}(x-y)\right]dr+2\int_0^t\left[p_{2t-2r}(0)-p_{2t-2r}(x-y)\right]dr \\ & -4\int_0^{s\wedge t}\left[p_{s+t-2r}(0)-p_{s+t-2r}(x-y)\right]dr.\end{aligned}$$

By change of variables $u = 2s - 2r$, $v = 2t - 2r$ and $w = s + t - 2r$ in the above corresponding integrals respectively and noticing that $s + t - 2(s \wedge t) = |t - s|$, we get

$$\begin{aligned} \mathbb{E}\left[\Box^2_{(s,x)} u(t,y)\right]^2 &= \int_0^{2s} [p_u(0) - p_u(x-y)]\, du + \int_0^{2t} [p_v(0) - p_v(x-y)]\, dv \\ &\quad -2 \int_{|s-t|}^{s+t} [p_w(0) - p_w(x-y)]\, dw \\ &= \left(\int_{s+t}^{2(s\vee t)} - \int_{2(s\wedge t)}^{s+t} + 2 \int_0^{|s-t|} \right) [p_r(0) - p_r(x-y)]\, dr. \end{aligned}$$

By Lemma 5.1, we see that

$$\begin{aligned} &\left(\int_{s+t}^{2(s\vee t)} - \int_{2(s\wedge t)}^{s+t} \right) [p_r(0) - p_r(x-y)]\, dr \\ &\qquad\qquad \le \frac{1}{\sqrt{2\pi}} \left(1 - e^{-\frac{(x-y)^2}{2(s+t)}} \right) \left(\sqrt{2s} + \sqrt{2t} - 2\sqrt{s+t} \right) \le 0. \end{aligned}$$

and

$$\int_0^{|s-t|} [p_r(0) - p_r(x-y)]\, dr \le c_\alpha |x-y|^{2\alpha} |s-t|^{1/2-\alpha}$$

for every α in $[0, 1/2]$. Thus

$$\mathbb{E}\left[\Box^2_{(s,x)} u(t,y)\right]^2 \le c_\alpha |x-y|^{2\alpha} |s-t|^{1/2-\alpha}.$$

An application of Theorem 4.3 immediately gives the desired result. ■

Remark 5.3 *Using the method in Remark 4.5, the above result implies the following estimate*

$$|u(s,x) - u(t,y)| \le C \left(|s-t|^{1/4} \sqrt{\log \frac{1}{|x||s-t|}} + |x-y|^{1/2} \sqrt{\log \frac{1}{|x-y||t|}} \right) \tag{5.4}$$

which is sharper than (5.2).

References

[1] Ayache, A.; Leger, S. and Pontier, M. Drap brownien fractionnaire. Potential Anal. 17 (2002), no. 1, 31-43.

[2] Ayache, A. and Xiao, Y. Asymptotic properties and Hausdorff dimensions of fractional Brownian sheets. J. Fourier Anal. Appl. 11 (2005), no. 4, 407-439.

[3] Fernique, X. Continuité des processus Gaussiens. C. R. Acad. Sci. Paris 258 (1964), 6058-6060.

[4] Fernique, X. Séries de distributions aléatoires indépendantes. C. R. Acad. Sci. Paris Sér. A-B 263 (1966), A674-A677.

[5] Fernique, X. Fonctions aléatoires gaussiennes, vecteurs aléatoires gaussiens. Université de Montréal, Centre de Recherches Mathématiques, Montreal, QC, 1997.

[6] Garsia, A. M.; Rodemich, E.; and Rumsey, H., Jr. A real variable lemma and the continuity of paths of some Gaussian processes. Indiana Univ. Math. J. 20 (1970/1971), 565-578.

[7] Ral'chenko, K. V. The two-parameter Garsia-Rodemich-Rumsey inequality and its application to fractional Brownian fields. Theory Probab. Math. Statist. No. 75 (2007), 167-178.

[8] Marcus, M. B.; Shepp, L. A. Continuity of Gaussian processes. Trans. Amer. Math. Soc. 151 (1970), 377-391.

[9] Xiao, Y. Sample path properties of anisotropic Gaussian random fields. (English summary) A minicourse on stochastic partial differential equations, 145212, Lecture Notes in Math., 1962, Springer, Berlin, 2009.